



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

PROBLEMS.

54. Proposed by E. W. MORRELL, Department of Mathematics, Montpelier Seminary, Montpelier, Vermont.

Transform $x^4 + y^4 + z^4 - 2y^2z^2 - 2z^2x^2 - 2x^2y^2$ into a product.
[Bowser's *Trigonometry*.]

55. Proposed by MARCUS BAKER, M. A., U. S. Geological Survey, Washington, D. C.

Two right triangles ABC and ABD are so placed as to have one side $x(=AB)$ in common. From P the intersection of their hypotenuses is drawn c perpendicular to x . Knowing the hypotenuses $a=39$ feet and $b=25$ feet and the per-

pendicular $c=12\frac{1}{2}$ feet, find x . Note this theorem $\frac{1}{m} + \frac{1}{n} = \frac{1}{c}$ or $\frac{1}{\sqrt{a^2 - x^2}}$.

$+ \frac{1}{\sqrt{b^2 - x^2}} = \frac{1}{c}$, where m and n are the altitudes of the two triangles, respectively. Also find locus of P . Discuss the case when the triangles are general (not right angled.)

[The same problem, in the form of "two poles" with ropes stretched from top of one to foot of other and the same data given, was contributed by H. C. Wilkes. Ed.]

GEOMETRY.

Conducted by B. F. FINKEL, Springfield, Mo. All contributions to this department should be sent to him.

SOLUTIONS OF PROBLEMS.

43. Proposed by J. F. W. SCHEFFER, Hagerstown, Maryland.

The consecutive sides of a quadrilateral are a, b, c, d . Supposing its diagonals to be equal, find them and also the area of the quadrilateral.

Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Mathematics, Inter State College, Texarkana, Texas.

Let $AB=a, BC=b, CD=c, DA=d, AC=DB=x, EF=y, GH=z$; also, let H, F, G, E, I, K be the middle points of AB, BC, CD, DA, AC, DB , respectively.

Then $EGFH, GKHI, FKEI$ are all parallelograms; but $IG=HK=\frac{1}{2}d, HI=KG=\frac{1}{2}b, EI=KF=\frac{1}{2}c, FI=EK=\frac{1}{2}a, EG=GF=EH=HE=\frac{1}{2}x$.

$$\therefore \frac{1}{4}(a^2 + c^2) = y^2 +$$

